

**MENG/SKUD GOED VOOR AANWENDING • MIX/SHAKE WELL BEFORE APPLICATION**

GROEP 2 MISSTOF  
GROUP 2 FERTILISER

WET / ACT NO. 36/1947  
REG. NO: B5060

LOT/BATCH #

VERVAARDIGINGSDATUM:  
DATE OF MANUFACTURE:



# FRUGVULT



**PLANTVOEDINGSTOF INHOUD**  
**PLANT NUTRITION CONTENT**

N	3.9	g/kg
P	5.5	g/kg
K	39.3	g/kg
Ca	4.0	g/kg
Mg	4.0	g/kg
S	8.3	g/kg

**GESKIK VIR**  
**BESPROEINGSTOEDIENING**  
**EN BLAARBESPUITING**

**IDEAL FOR FERTIGATION**  
**AND FOLIAR SPRAY**

**VERSPREI DEUR / DISTRIBUTED BY:**



**VERVAARDIG DEUR:**  
**MANUFACTURED BY:**



CK 1999/066144/23  
BTW/VAT: 4170199014

**POSBUS/P.O. BOX 341**  
**VANDERBIJLPARK 1900**

**SUID-AFRIKA / SOUTH AFRICA**

**T: +27 (0)16 987 2862**

**E: info@nutracor.co.za**

**VERSEËL ONGEBRUIKTE PRODUK**  
**SEAL UNUSED PRODUCT**

**VOLUME 5, 10 & 20L**

**HOU WEG VAN VOEDSEL**  
**KEEP AWAY FROM FOODS**



# FRUGVULT

GEWAS CROP	BESPROEING FERTIGATION	OPMERKINGS: REMARKS:
GROENTE VEGETABLES	250-500ml /100L water	Begin toedienings net na set en dien toe elke 2 weke tydens vrugontwikkeling van tamaties en rissies. Start applications after set and repeat every 2 weeks during fruit growth period of tomatoes and peppers.
VRUGTEBOME/ KERN EN STEEN FRUIT TREES/ POME AND STONE	250-500ml /100L water	3 - 4 toedienings vanaf 100% blomblaarval gedurende vrugontwikkeling tot week voor oes. 3 - 4 applications starting at 100% petal fall during fruit growth period till week before harvest.
SITRUS CITRUS	250-500ml /100L water	2 - 3 toedienings vanaf 100% blomblaarval elke 2 tot 3 weke. 2 - 3 applications starting at 100% petal fall every 2 to 3 weeks.
TAFELDRUIWE EN WINGERDE TABLE GRAPES AND VINEYARDS	250-500ml /100L water	3 - 4 toedienings net na set, ertjiekorrel, en elke twee weke tot week voor deurslaan. 3 - 4 applications after set, pea size, and every two weeks until week before veraison.
SUBTROPIESE VRUGTE EN NEUTE SUBTROPICAL FRUITS AND NUTS	250-500ml /100L water	2 - 3 toedienings elke 3 tot 4 weke tydens vrugontwikkeling vanaf set. 2 - 3 applications every 3 to 4 weeks during fruit growth period starting after fruit set.

\*KONTAK U VERSPREIDER, KONSULTANT OF VERVAARDIGER VIR 'N SPESIFIEKE GEWASPROGRAM EN PRODUKAANBEVELINGS.

\*CONTACT YOUR DISTRIBUTOR, CONSULTANT OR THE MANUFACTURER FOR A SPECIFIC CROP PROGRAM AND PRODUCT RECOMMENDATIONS.

Die vervaardiger van topGroei produkte is nie verantwoordelik/aanspreeklik vir enige wanaanwending van die produkreeks.

The manufacturer of topGroei products is not responsible for any erroneous application of the product range.

**MENG/SKUD GOED VOOR AANWENDING • MIX/SHAKE WELL BEFORE APPLICATION**

VERSEËL ONGEBRUIKTE PRODUK • SEAL UNUSED PRODUCT

HOU WEG VAN VOEDSEL • KEEP AWAY FROM FOODS